


AST-101



Solar System: origin and exploration

Luís Anchordoquí

Origin of the Solar System



What any theory must address?

- Why do the orbits of the planets lie in the same plane ?
- Why do the eight planets and asteroids all orbit the Sun with prograde revolution?
- Why does the Sun rotate in this prograde sense?
- Why are the orbits of the planets nearly circular?
- Why do six of the eight planets have prograde rotation?
(With the exception of Venus and Uranus)
- Why does Bode's Rule describe the nearly regular spacing of planetary orbits? (Each planet has approximately twice the distance to the Sun as its nearest inner neighbor)

What any theory must address?


Questions of Planetary Composition

- Why do the planets vary systematically in composition and density, with the innermost planets tending toward high density and heavy elements, and the outermost planets tending toward low density and substances of low melting point?

Gross Properties of the Planets

Planet	Distance AU	Diameter 10^3 km	Mass (Earth = 1)	Density (gm/cm^3)	Orbital Inclination
Mercury	0.39	4.9	0.055	5.4	7.0°
Venus	0.72	12.1	0.82	5.25	3.6°
Earth	1.00	12.7	1.00	5.52	--
Mars	1.52	6.8	0.11	3.93	1.9°
Jupiter	5.20	143	318	1.33	1.3°
Saturn	9.54	120	95	0.71	2.5°
Uranus	19.2	51	15	1.27	0.8°
Neptune	30.1	50	17	1.70	1.8°
Pluto	39.4	2.4	0.03	1.99	17.2°

Encounter Hypothesis



Encounter Hypothesis

(Buffon 1745)

1. A “near miss” encounter occurs between the Sun and a passing star
2. Material is pulled from the stellar surfaces by tidal forces
3. Material cools and condenses to form planetary bodies

Principal Shortcoming: This Requires a highly unlikely event

Calculation ➡ Collision rate is proportional to

- Relative speed
- Collision Cross-Section (How close is “close”?)
- Density of stars

This gives a typical collisional interval of 10^{20} years!

Predictive Successes ➡

Direct, coplanar, low eccentricity orbits

Common planetary ages (but an older Sun)

But it Doesn't Work ➡



Ejected gas escapes before cooling

Conservation of Energy

Energy = Kinetic Energy + Potential Energy

$$E = KE + PE = \frac{1}{2}mv^2 + mgy$$

$v = 0$



Energy buckets (right) show how energy moves from all potential to all kinetic

Ejected Gas Escapes before Cooling

$$\text{Potential Energy} = \frac{GMm}{r}$$

$$\text{Kinetic Energy} = \frac{1}{2} mv^2$$


Initial Kinetic Energy = Potential Energy

$$\frac{1}{2}mv^2 = \frac{GMm}{r} \quad \Rightarrow \quad v_{\text{escape}}^2 = \frac{2GM}{r}$$


$$v_{\text{thermal}}^2 = 3kT/m \quad k \Rightarrow \text{Boltzmann constant}$$

For high temperature $T \Rightarrow v_{\text{escape}}^2 \ll v_{\text{thermal}}^2$


Nebular Hypothesis




The solar nebula contracts



As the nebula shrinks,
its motion causes it to flatten



The nebula is a disk of matter
with a concentration near the center





Formation of the protosun
Solid particles condense as the nebula cools,
giving rise to the planetesimals,
which are the building blocks of the planets

Nebular Hypothesis & Protoplanets


- The Sun forms from a collapsing cloud of **cold** interstellar gas and dust
- The material forms a proto-Sun surrounded by a **cool** gas and dust disk
- Small particles form and grow in the disc by **collisional accretion**
- Larger bodies (**planetesimals**) accrete rapidly with the aid of gravity
- Planetesimals grow by accretion of gas, dust, and other planetesimals
 - gradually clearing the disk of remaining material.
 - maintained in gaseous states by collisional heating
- Final states of planetesimals growth form protoplanets
 - surfaces cooling and solidifying as accretion slows
 - atmospheres dissipate at a decreasing rate
- The protoplanets evolve with time to become the present-day planets
 - differentiation between inner and outer planets

Note: Satellites are formed by accretion in disks about protoplanets or acquired by later capture or result from collisions





Inner planets have all of their volatile elements "boiled off" by the heat of the Sun (hydrogen, lithium, helium)



Jovian planets do not receive enough solar radiation for this to happen so the lighter elements remain and accumulate over time

Nebular Hypothesis

Some History

- ☑ Descartes 1644 ➡ Eddies and whirlpools in a turbulent disk
- ☑ Kant 1755 ➡ Planet formation in a rotating Keplerian disk
- ☑ Laplace 1796 ➡ A disk with “tidal rings”

Some Virtues

- ☑ Cool early environment allows formation of larger solid bodies
- ☑ Planet and star (Sun) formation are part of a single process ➡ accounts for gross orbital properties
- ☑ Accounts for composition similarities and differences (Jovian/Terrestrial)


Current Status

- ☑ Very great explanatory and predictive power > Eminently testable
- ☑ No fundamental issues or problems outstanding
- ☑ Is a successful model for star formation as well as stellar systems


(Stars are now known and observed to form from cool interstellar clouds)

- ☑ No viable competing model has been forthcoming

The sun still contains most of the material of the original solar nebula



Its internal nuclear reactions have modified the material at the sun's core
However, the surface layers, which have not mixed with the core in its present state, have quite accurately preserved the original nebular composition



- The solar wind is a flow of particles that comes off the sun at about one million miles per hour and travels throughout the solar system
- At the outer limits of the solar system, the solar wind finally begins to lose out and its speed drops below the speed of sound (relative to the interstellar medium), resulting in a roughly spherical shell known as the termination shock front
- Heliopause is the boundary where the interstellar medium and solar wind pressures balance
- The bow shock is typically the boundary between the stellar wind and the interstellar medium

*Interstellar
Winds*




*Outer boundary
of Heliosphere*

Sun

*Subsonic
Flow*

*Shock terminating
the region of supersonic
solar wind*



Exploration of the Solar System







Apolo 11 ➤ July 16, 1969 @13:32 UTC




Apollo 11 Lunar Module Eagle, in a landing configuration was photographed in lunar orbit from the Command and Service Module Columbia

Neil Armstrong ➡ first step on the Moon
July 20, 1969 @20:17 UTC









An aerial photograph of the Apollo 11 Lunar Module (LM) floating in the ocean. The LM is yellow and conical, with three white spherical flotation devices attached to its top. It is surrounded by a large orange life raft. To the right of the LM, a smaller orange life raft contains three astronauts in green suits and red life preservers. The water is a deep blue with small waves.


Apollo 11 Comes Home ➡ Armstrong, Aldrin, and Collins
splash down on July 24, 1969 @ 16:50 UTC

Apollo 11 timeline










Jim Irwin sets up the first Lunar Rover Vehicle during Apollo 15 in 1971




Mariner 10 was an American robotic space probe launched by NASA on November 3, 1973 to fly by the planets Mercury and Venus

Voyager the Grand Tour and beyond





Uranus and Neptune with Voyager 2, Picture from Neptuno was taken at a range of 4.4 million miles from the planet, 4 days and 20 hours before closest approach




The Soviet Mars 3 achieved the first soft landing on Mars in 1971, but its instruments stopped working after only 20 seconds on the surface

The Twin Vikings landed on Mars in 1976

Viking 2 returned data until 1980, and contact with Viking 1 was lost in late 1982


Carl Sagan next to Viking lander mockup in Death Valley, California



The Mars Orbiter Camera was in orbit around Mars between 1997 and 2006
(MOC)

The largest volcano in the solar system

MOC obtained this spectacular wide-angle view of Olympus Mons on Mars Global Surveyor's 263rd orbit (April 25 1998)



Giant volcano on Mars ➡ three times as high as Everest and as wide as France
↓
8.8km


NASA's Spirit and Opportunity Mars rovers landed on the Red Planet on Jan. 3 and 24, 2004, respectively



This image shows a view Opportunity captured of its own shadow on July 26 of that year, the 180th Martian day of its mission


2003-06-10 18:35

Spirit



2003-06-17 06:35

Spirit




32.3km/s

94,688,685km

(88.4 - 168.7)

2003-06-17 18:35

Spirit




34
2003-06-23 18:35

Spirit

32.1km/s


87,017,351km

(68.4 - 168.7)




2003-06-26 06:35

Spirit



34 2003-07-04 06:35

Spirit




34
2003-07-12 18:35

Spirit

31.4km/s

65,645,494km

(88.4 - 168.7)




34 2003-07-23 06:35

Spirit

30.9km/s

56,935,221km

(88.4 - 168.7)




34 2003-08-03 18:35

Spirit

30.2km/s


47,699,556km

(88.4 - 168.7)




2003-08-11 06:35

Spirit




34
2003-08-20 06:35

Spirit




2003-08-28 06:35

Spirit




2003-09-07 18:35

Spirit




34 2003-09-20 18:35

Spirit



2003-10-07 06:35

Spirit




26.0km/s

19,607,746km

(88.4 - 168.7)

2003-10-17 06:35

Spirit




25.4km/s

17,373,021km

(88.4 - 168.7)

2003-10-24 18:35

Spirit




25.0km/s

15,787,266km

(88.4 - 168.7)


2003-11-02 06:35

Spirit




2003-11-11 18:35

Spirit



2003-11-17 18:35

Spirit




23.8km/s

10,774,221km

(88.4 - 168.7)


2003-11-27 06:35

Spirit




34 2003-12-01 06:35

Spirit



34 2003-12-09 06:35

Spirit




23.0km/s

6,036,040km

(88.4 - 168.7)

34 2003-12-14 06:35

Spirit




22.8km/s

4,894,585km

(88.4 - 168.7)


34 2003-12-21 06:35

Spirit




2003-12-24 18:35

Spirit



34 2003-12-29 06:35

Spirit




22.4km/s

1,409,675km

(88.4 - 168.7)

34 2003-12-31 18:35

Spirit




22.3km/s

822,068km

(88.4 - 168.7)

34 2004-01-04 04:20

Spirit



20.4km/s

3,523km

(88.4 - 168.7)

Spirit and Opportunity BY THE NUMBERS




6 YEARS
lifespan

128,000
raw images

4.8 MILES
traveled

30 DEGREES
steepest slope



10 YEARS
lifespan

187,000
raw images


24 MILES
traveled

31 DEGREES
steepest slope




Image captured by the Spirit Rover on May 19, 2005


Curiosity Mars Science Laboratory




Curiosity was launched from Cape Canaveral on November 26, 2011 @ 15:02 UTC
and landed on Aeolis Palus inside Gale crater on Mars on August 6, 2012 @ 05:17 UTC




**Curiosity sent back a striking selfie captured using
the camera on its robotic arm**



Galileo orbited Jupiter from 1995 to 2003, and conducted flybys of several of its moons...




...including this one: Europa



This image of Jupiter was actually taken by the Cassini spacecraft en route to Saturn



Image of Saturn was taken by the Cassini spacecraft from a distance of 177 million miles (285 km) as it approached the planet. The little dot to Saturn's upper left is Titan, whose brightness has been enhanced relative to Saturn



43

Launched in 1997, NASA's Cassini space probe arrived in Saturn orbit in 2004
On July 19 of this year the spacecraft took an extraordinary image of our
home planet and moon as seen from nearly 900 miles away

The long road to a Pluto mission



1977
Voyager 1
722 kilograms
\$1.7 billion*



1990
Pluto 350
350 kilograms
\$543 million



1991
Mariner Mark II
2,000 kilograms
\$3.2 billion



1991
Pluto Fast Flyby
140 kilograms
\$2.1 billion



1995
Pluto Express
Pluto Kuiper Express
175 kilograms
\$1.4 billion



2001
New Horizons
385 kilograms
\$646 million

All costs converted to 2014 dollars.
*Total cost of Voyager program through 1989



Pluto as seen by New Horizons



Charon as seen by New Horizons




Comet Tempel 1 image was taken 67 seconds after it obliterated Deep Impact's impactor spacecraft

As comets travel close to the Sun, the Sun's heat begins to vaporize the ices and causes them to form a fuzzy, luminous area of vaporized gas around the nucleus of the comet known as a coma

Outside the coma is a layer of hydrogen gas called a hydrogen halo which extends up to 10^{10} meters in diameter

The solar wind then blows these gases and dust particles away from the direction of the Sun causing two tails to form




These tails always point away from the Sun as the comet travels around it

One tail is called the ion tail and is made up of gases which have been broken apart into charged molecules and ions by the radiation from the Sun

Since the most common ion, CO^+ scatters the blue light better than red light, to observers, this ion tail often appears blue

The other tail is called a dust tail and normally appears white



That is why the dust tail is usually curved rather than straight, and does not point directly away from the Sun, because it is also influenced by the motion of the comet

The tails of the comet can be extremely large and may extend a distance of up to 1 AU (the distance between the Earth and the Sun)!




Both tails can be seen in the image of Comet Hale-Bopp

'Oumuamua


- First object ever detected to have visited our solar system from another star
- It was first detected via telescope in 2017 by Robert Weryk as it passed through our solar system and sling-shotted past the sun at nearly 200,000 miles per hour
- Almost as soon as the object was discovered, it was on its way out, passing the orbits of Mars, then Jupiter in 2018, and then Saturn in 2019



A Mission to 'Oumuamua?



- Large uncertainty in its current position
- At its current distance from the Sun, 'Oumuamua is more than a million times fainter than it was near Earth and any targeting spacecraft will need to be equipped with a large telescope to find 'Oumuamua



‘Oumuamua (pronounced oh-MOO-ah-MOO-ah) means
“a messenger from afar arriving first” in Hawaiian

The object appears to have come from roughly the direction of
Vega in the constellation Lyra and is now far beyond the orbit of
Neptune heading toward the constellation Pegasus

Vega is a star relatively close at only 25 light-years from the Sun
















































































What planet has the highest mountain in the Solar System?

A. Venus

B. Earth

C. Mars

What planet has the highest mountain in the Solar System?

A. Venus

B. Earth

C. Mars

What was the first space probe to leave the Solar System?

A. The Hubble Telescope

B. Voyager 1

C. Voyager 2

D. Halley Armada

What was the first space probe to leave the Solar System?

A. The Hubble Telescope

B. Voyager 1

C. Voyager 2

D. Halley Armada

How long would a one-way mission from Earth to Mars take?

- A. 7 days
- B. 7 weeks
- C. 7 months
- D. 7 years

How long would a one-way mission from Earth to Mars take?

A. 7 days

B. 7 weeks

C. 7 months

D. 7 years

Comets have two tails. One is a dust tail comprises of particles from its surface

What is the other tail called..?

A. Ion

B. Stream

C. Double tail

Comets have two tails. One is a dust tail comprises of particles from its surface

What is the other tail called..?

A. Ion

B. Stream

C. Double tail

The nucleus of a comet is surrounded by..?

A. A tail

B. A coma

C. A full stop

The nucleus of a comet is surrounded by..?

A. A tail

B. A coma

C. A full stop

How old is the Solar System

A. 4.6 billion years

B. 1 million years

C. 100,000 years

D. 10,000 years

How old is the Solar System

A. 4.6 billion years

B. 1 million years

C. 100,000 years

D. 10,000 years

What is the most widely accepted theory for the formation of the solar system?

- A. Nebular hypothesis**
- B. Ptolemaic Model**
- C. Encounter hypothesis**

What is the most widely accepted theory for the formation of the solar system?

A. Nebular hypothesis

B. Ptolemaic Model

C. Encounter hypothesis

Who was the first man to walk on the Moon?

A. Buzz Aldrin

B. Neil Armstrong

C. Michael Collins

Who was the first man to walk on the Moon?

A. Buzz Aldrin

B. Neil Armstrong

C. Michael Collins

What was the name of the mission?

A. Apollo 11

B. Starbuck 11

C. Halo 11

What was the name of the mission?

A. Apollo 11

B. Starbuck 11

C. Halo 11

What year did a man first walk on the Moon?

A. 1966

B. 1967

C. 1968

D. 1969

What year did a man first walk on the Moon?

A. 1966

B. 1967

C. 1968

D. 1969

What was the name of the rocket they launched in?

A. Mars V

B. Jupiter V

C. Saturn V

What was the name of the rocket they launched in?

A. Mars V

B. Jupiter V

C. Saturn V

How long did the mission take?

A. 6 days

B. 8 days

C. 10 days

How long did the mission take?

A. 6 days

B. 8 days

C. 10 days

‘Oumuamua was discovered by Robert Weryk using the Pan-STARRS telescope at

- A. Haleakalā Observatory, Hawaii
- B. Arecibo Observatory, Puerto Rico
- C. Argentine Institute of Radio Astronomy, Berazategui, Argentina
- D. Arcetri Observatory, Florence, Italy
- E. Astronomy Tower of the Sorbonne, Paris, France

‘Oumuamua was discovered by Robert Weryk using the Pan-STARRS telescope at

- A. Haleakalā Observatory, Hawaii**
- B. Arecibo Observatory, Puerto Rico**
- C. Argentine Institute of Radio Astronomy, Berazategui, Argentina**
- D. Arcetri Observatory, Florence, Italy**
- E. Astronomy Tower of the Sorbonne, Paris, France**

QUERY 9

Determine the escape velocity of Earth, Mars, and the Moon

QUERY 9

Determine the escape velocity of Earth, Mars, and the Moon

$$v_{\text{escape,Earth}} = \sqrt{2GM/R} = \sqrt{2 \cdot 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2 \cdot 5.9 \times 10^{24} \text{ kg} / (6.4 \times 10^6 \text{ m})} = 1.1 \times 10^4 \text{ m/s} = 11 \text{ km/s}$$

$$v_{\text{escape,Mars}} = \sqrt{2GM/R} = \sqrt{2 \cdot 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2 \cdot 6.5 \times 10^{23} \text{ kg} / (3.4 \times 10^6 \text{ m})} = 5.0 \times 10^3 \text{ m/s} = 5 \text{ km/s}$$

$$v_{\text{escape,Moon}} = \sqrt{2GM/R} = \sqrt{2 \cdot 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2 \cdot 7.2 \times 10^{22} \text{ kg} / (1.7 \times 10^6 \text{ m})} = 2.4 \times 10^3 \text{ m/s} = 2.4 \text{ km/s}$$

QUERY 10

The escape speed from Earth is 11 km/s, so a gas molecule travelling away from Earth near the outer boundary of the Earth atmosphere would, at this speed, be able to escape from the Earth gravitational field. Actually, if the thermal speed of a gas is greater than about 15 to 20% of the escape speed of a planet, virtually all of the molecules of that gas will escape the atmosphere of the planet.

At what temperature is the thermal speed of (i) oxygen molecules and (ii) helium molecules equal to 1.65 km/s

(iii) Explain why our atmosphere contains oxygen but not helium

[Hint: The masses of one molecule of O_2 (molecular mass = 32 u) and

He (molecular mass = 4 u) are $m(O_2) = 5.3 \times 10^{-26}$ kg

$m(He) = 6.6 \times 10^{-27}$ kg where $1 \text{ u} = 1.66 \times 10^{-27}$ kg

QUERY 10

(i)

$$T_{\text{O}_2} = \frac{(0.15 v_{\text{escape}})^2 m}{3k} = \frac{32 \cdot 1.66 \times 10^{-27} \text{ kg} \cdot (0.15 \cdot 1.1 \times 10^4 \text{ m/s})^2}{3 \cdot 1.38 \times 10^{-23} \text{ J/K}} = 3,708 \text{ K}$$

(ii)

$$T_{\text{He}} = \frac{(0.15 v_{\text{escape}})^2 m}{3k} = \frac{4 \cdot 1.66 \times 10^{-27} \text{ kg} \cdot (0.15 \cdot 1.1 \times 10^4 \text{ m/s})^2}{3 \cdot 1.38 \times 10^{-23} \text{ J/K}} = 46 \text{ K}$$

(iii) Because the **escape temperature** is so high for oxygen, very few oxygen molecules ever escape the atmosphere

But helium, with one-eighth the mass, can escape at much lower temperature

The average surface temperature on Earth is approximately 15°C

QUERY 11

Compute the temperature for which the thermal speed of O_2 is equal to 15% of the escape speed at the surface of the Moon. How does this account for the absence of an atmosphere on the Moon?


QUERY 11

Compute the temperature for which the thermal speed of O_2 is equal to 20% of the escape speed at the surface of the Moon. How does this account for the absence of an atmosphere on the Moon?

$$T_{\text{O}_2} = \frac{(0.15 v_{\text{escape}})^2 m}{3k} = \frac{32 \cdot 1.66 \times 10^{-27} \text{ kg} \cdot (0.15 \cdot 2.3 \times 10^3 \text{ m/s})^2}{3 \cdot 1.38 \times 10^{-23} \text{ J/K}} = 162 \text{ K}$$

The mean surface temperature of the Moon is 106°C during the day and -183°C during night. The temperature during the day is hot enough that oxygen (and other gases) would have escaped during the time since the formation of the moon to the present. Moreover, the temperature during the moon formation could have been approximately 1000 K.

Pale Blue Dot



Consider again that blue dot. That's here. That's home. That's us.

On it everyone you love, everyone you know, everyone you ever heard of, every human being who ever was, lived out their lives.

The aggregate of our joy and suffering, thousands of confident religions, ideologies, and economic doctrines, every hunter and forager, every hero and coward, every creator and destroyer of civilization, every king and peasant, every young couple in love, every mother and father, hopeful child, inventor and explorer, every teacher of morals, every corrupt politician, every "superstar," every "supreme leader," every saint and sinner in the history of our species lived there – on a mote of dust suspended in a sunbeam

The Earth is a very small stage in a vast cosmic arena. Think of the rivers of blood spilled by all those generals and emperors so that, in glory and triumph, they could become the momentary masters of a fraction of a dot. Think of the endless cruelties visited by the inhabitants of one corner of this pixel on the scarcely distinguishable inhabitants of some other corner, how frequent their misunderstandings, how eager they are to kill one another, how fervent their hatreds.

Our posturings, our imagined self-importance, the delusion that we have some privileged position in the Universe, are challenged by this point of pale light. Our planet is a lonely speck in the great enveloping cosmic dark. In our obscurity, in all this vastness, there is no hint that help will come from elsewhere to save us from ourselves.

The Earth is the only world known so far to harbor life. There is nowhere else, at least in the near future, to which our species could migrate. Visit, yes. Settle, not yet. Like it or not, for the moment the Earth is where we make our stand.

It has been said that astronomy is a humbling and character-building experience. There is perhaps no better demonstration of the folly of human conceits than this distant image of our tiny world. To me, it underscores our responsibility to deal more kindly with one another, and to preserve and cherish the pale blue dot, the only home we've ever known.

Carl Sagan